arXiv:math/0102013v3 [math.AG] 25 Dec 2003

CHARACTERISTIC NUMBERS OF A HOMOGENEOUS SPACE

LORING W. TU

Abstract. Let G be a compact connected Lie group with maximal torus T, and H a closed subgroup containing T. We work out the Atiyah-Bott-Berline-Vergne localization formula for the homogeneous space G/H under the natural action of the maximal torus T. The computation gives explicit formulas for the ordinary and equivariant characteristic numbers of a homogeneous space.

Let G be a compact connected Lie group with maximal torus T, and H a closed subgroup of G containing T. The quotient space G/H includes examples such as Grassmannians and flag manifolds. The torus T acts on G/H by left multiplication. In this note we apply the Atiyah-Bott-Berline-Vergne localization formula to calculate the equivariant characteristic numbers of G/H under this torus action (Section 13).

As it well known, this yields the ordinary characteristic numbers of the homogeneous space G/H (Th. 14). However, our main interest is to derive a restriction formula (Th. 8) and an Euler class formula (Th. 12) in the equivariant cohomology of G/H, for it turns out that these formulas can be used to calculate the Gysin homomorphism in the ordinary cohomology of a flag bundle. The application to the Gysin homomorphism is an instance of a general principle relating the push-forward in ordinary cohomology to the localization formula in equivariant cohomology. As this more general principle is independent of homogeneous spaces, we treat it in a separate article [15].

In [13] a relative localization formula is used to calculate the Gysin map of an equivariant map. The restriction formula (Th. 8) and the Euler class formula (Th. 12) of the present article enter essentially in a calculation that allows us to recover a formula of Akyildiz and Carrell [2] for the Gysin map of a map between flag manifolds.

Using techniques of symplectic quotients, Shaun Martin ([12], Prop. 7.2) has obtained a formula for the characteristic numbers of a Grassmannian similar to our Prop. 15.

To simplify the exposition, we discuss first the homogeneous space G/T. The general case G/H follows by suitable modifications.

Although we consider cohomology with real coefficients, everything goes through with rational coefficients. To minimize the pre-requisites and to establish the notations, in the first few sections we recall some results on associated bundles, the characteristic map, the actions of the Weyl group, and equivariant cohomology.

I am grateful to Raoul Bott for suggesting the problem and for many helpful discussions.

1. Associated bundles and the characteristic map

Suppose a torus T of dimension ℓ acts freely on the right on a topological space X so that $X \longrightarrow X/T$ is a principal T-bundle. A *character* of T is a multiplicative homomorphism of T into $\mathbb{C}^*$. Let $\hat{T}$ be the group of characters of T, with the multiplication of characters written additively: for $\alpha, \beta \in \hat{T}$ and $t \in T$,

$$t^{\alpha+\beta} := t^\alpha t^\beta = \alpha(t)\beta(t).$$

Denote by $\mathbb{C}_\gamma$ the complex vector space $\mathbb{C}$ with an action of T given by the character $\gamma : T \longrightarrow \mathbb{C}^*$.

By the mixing construction, a character γ on T associates to the principal bundle $X \longrightarrow X/T$ a complex line bundle $L(X/T, \gamma)$ over X/T:

$$L(X/T, \gamma) := X \times_T \mathbb{C}_\gamma := (X \times \mathbb{C}_\gamma)/T,$$

Date: December 25, 2003.

Key words and phrases. Atiyah-Bott-Berline-Vergne localization formula, equivariant cohomology, equivariant characteristic numbers, homogeneous spaces.

2000 Mathematics Subject Classification. Primary: 57R20, 14M17; Secondary: 55N25, 14C17.

The author acknowledges the hospitality and support over a period of several years of the Institut Henri Poincaré, the École Normale Supérieure, the Institut de Mathématiques de Jussieu, and the Ministère de l'Enseignement et de la Recherche, France.

where T acts on $X \times \mathbb{C}_\gamma$ by

$$(x, v)t = (xt, \gamma(t^{-1})v).$$

The equivalence class of (x, v) is denoted $[x, v]$.

The first Chern class of an associated line bundle defines a homomorphism of abelian groups

$$\begin{aligned} c : \hat{T} &\longrightarrow H^2(X/T) \\ \gamma &\mapsto c_1(L(X/T, \gamma)). \end{aligned} \tag{1}$$

Let $\operatorname{Sym}(\hat{T})$ be the symmetric algebra with real coefficients generated by the additive group $\hat{T}$. The group homomorphism (1) extends to a ring homomorphism

$$c : \operatorname{Sym}(\hat{T}) \longrightarrow H^*(X/T),$$

called the *characteristic map* of X/T, sometimes denoted $c_{X/T}$.

We apply the construction of the associated bundle of a character $\gamma \in \hat{T}$ to two situations:

1) The classifying space BT. Applied to the universal bundle $ET \longrightarrow BT$, this construction yields line bundles
$$S_\gamma := L(ET/T, \gamma)$$
over BT and cohomology classes $c(\gamma) = c_1(S_\gamma)$ in $H^2(BT)$. The characteristic map
$$c : \operatorname{Sym}(\hat{T}) \longrightarrow H^*(BT)$$
is a ring isomorphism, since both are polynomial rings in ℓ generators and the generators correspond. If $\chi_1, \dots, \chi_\ell$ is a basis for the character group $\hat{T}$ and $u_i = c_1(S_{\chi_i})$, then $\operatorname{Sym}(\hat{T})$ is the polynomial ring $\mathbb{R}[\chi_1, \dots, \chi_\ell]$, and
$$H^*(BT) \simeq \mathbb{R}[u_1, \dots, u_\ell].$$
2) The homogeneous space G/T. Applied to the principal T-bundle $G \longrightarrow G/T$, this construction yields line bundles
$$L_\gamma := L(G/T, \gamma)$$
over G/T.

2. The actions of the Weyl group

The Weyl group of a maximal torus T in the compact connected Lie group G is $W = N_G(T)/T$, where $N_G(T)$ is the normalizer of T in G. The Weyl group is a finite reflection group.

We use w to denote both an element of W and a lift of the element to the normalizer $N_G(T)$. The Weyl group W acts on the character group $\hat{T}$ of T by

$$(w \cdot \gamma)(t) = \gamma(w^{-1}tw).$$

This action induces an action on $\operatorname{Sym}(\hat{T})$ as ring isomorphisms. Let R be the polynomial ring

$$R = \operatorname{Sym}(\hat{T}) = \mathbb{R}[\chi_1, \dots, \chi_\ell]$$

and R^W the subring of W-invariant polynomials.

If the Lie group G acts on the right on a space X, then the Weyl group W acts on the right on X/T by

$$r_w(xT) = (xT)w = xwT.$$

This action of W on X/T induces by the pullback an action of W on the line bundles over X/T and also on the cohomology ring $H^*(X/T)$: if L is a line bundle on X/T and $a \in H^*(X/T)$, then

$$w \cdot L = r_w^* L, \qquad w \cdot a = r_w^* a.$$

Proposition 1. *The action of the Weyl group W on the associated line bundles over X/T is compatible with its action on the characters of T; more precisely, for $w \in W$ and $\gamma \in \hat{T}$,*

$$r_w^* L(X/T, \gamma) \simeq L(X/T, w \cdot \gamma).$$

Indication of proof. Define $\varphi : r_w^* L(X/T,\gamma) \longrightarrow L(X/T, w\cdot\gamma)$ by

$$\varphi(xT, [xw, v]) = [x, v].$$

If we use a different representative xt to represent xT, then

$$(xT, [xw, v]) = (xtT, [xtw, v'])$$

and it is easily verified that $v' = \gamma(w^{-1}t^{-1}w)v$. Hence,

$$\begin{aligned}\varphi(xtT, [xtw, v']) &= [xt, v'] = [xt, (w\cdot\gamma)(t^{-1})v] \\ &= [x, v].\end{aligned}$$

This shows that φ is well defined. It has the obvious inverse map

$$\psi([x, v]) = (xT, [xw, v]).$$

□

Corollary 1.1. *The characteristic map* $c : \mathrm{Sym}(\hat{T}) \longrightarrow H^*(X/T)$ *is* W*-equivariant.*

Proof. Writing L_γ instead of $L(X/T,\gamma)$, it follows from the proposition that

$$r_w^* c(\gamma) = r_w^* c_1(L_\gamma) = c_1(r_w^* L_\gamma) = c_1(L_{w\cdot\gamma}) = c(w\cdot\gamma).$$

□

Corollary 1.2. *For* γ *a character on* T *and* w *an element of the Weyl group* W,

$$w \cdot c_1(S_\gamma) = c_1(S_{w\cdot\gamma}).$$

3. Homotopy quotients and equivariant cohomology

Suppose a group G acts on the left on a space M, and $EG \longrightarrow BG$ is the universal principal G-bundle. Since G acts freely on EG, the diagonal action of G on $EG \times M$,

$$(e, x)g = (eg, g^{-1}x),$$

is also free. The space $M_G := EG \times_G M := (EG \times M)/G$ is called the *homotopy quotient of* M *by* G, and its cohomology $H^*(M_G)$ the *equivariant cohomology of* M *under the* G*-action*, denoted $H_G^*(M)$. For the basics of equivariant cohomology, we refer to [7] or [10].

A G-equivariant vector bundle $E \longrightarrow M$ induces a vector bundle $E_G \longrightarrow M_G$. An *equivariant characteristic class* $c^G(E)$ of $E \longrightarrow M$ is defined to be the corresponding ordinary characteristic class $c(E_G)$ of $E_G \longrightarrow M_G$.

By the definition of the homotopy quotient, $M_G \longrightarrow BG$ is a fiber bundle with fiber M. Let $i : M \longrightarrow M_G$ be the inclusion map as a fiber. We say that a cohomology class $\tilde{\alpha} \in H_G^*(M)$ is an *equivariant extension* of $\alpha \in H^*(M)$ if $i^*\tilde{\alpha} = \alpha$. For example, if $E \longrightarrow M$ is a G-equivariant vector bundle, then any characteristic class $c(E)$ has an equivariant extension $c^G(E)$, since

$$i^* c^G(E) = i^* c(E_G) = c(i^* E_G) = c(E).$$

Now suppose G is a compact connected Lie group with maximal torus T. Let γ be a character of T. The torus acts by left multiplication on the associated line bundle $L_\gamma := L(G/T, \gamma) := G \times_T \mathbb{C}_\gamma$,

$$t[g, v] = [tg, v],$$

making $L_\gamma \longrightarrow G/T$ into a T-equivariant complex line bundle. Taking the homotopy quotients yields a complex line bundle $K_\gamma := (L_\gamma)_T \longrightarrow (G/T)_T$. Fix a basis $\chi_1, \dots, \chi_\ell$ for the characters of T and let $y_i = c_1(K_{\chi_i}) \in H_T^2(G/T)$. These are equivariant extensions of the cohomology classes $z_i := c_1(L_{\chi_i}) \in H^2(G/T)$.

Let $\pi : G/T \longrightarrow \mathrm{pt}$ be the constant map. Then

$$H_T^*(\{\mathrm{pt}\}) = H^*(BT) = \mathbb{R}[u_1, \dots, u_\ell], \qquad u_i = c_1(S_{\chi_i}). \tag{2}$$

It follows from ([8], Prop. 1(iii)) that the equivariant cohomology ring of G/T under the T-action is

$$H_T^*(G/T) = \mathbb{R}[u_1, \dots, u_\ell, y_1, \dots, y_\ell]/J, \tag{3}$$

where u_i stands for $\pi^* u_i$ and J is the ideal generated by $p(y) - p(u)$, as p ranges over all W-invariant polynomials of positive degree in ℓ variables, i.e., $p \in R_+^W$.

4. Restriction to a fixed point in G/T

For the action of the torus T on G/T by left multiplication, it is easily verified that the fixed point set is the Weyl group $W = N_G(T)/T$, considered as a subset of G/T. If $w \in W$ is a fixed point, let $i_w : \{w\} \hookrightarrow G/T$ be the inclusion map. Since i_w is T-equivariant, it induces a homomorphism in equivariant cohomology

$$i_w^* : H_T^*(G/T) \longrightarrow H_T^*(\{w\}).$$

Proposition 2. *With the identification*

$$H_T^*(G/T) = \mathbb{R}[u_1, \ldots, u_\ell, y_1, \ldots, y_\ell]/J$$

as in Eq. (3) *and*

$$H_T^*(\{w\}) \simeq H^*(BT) = \mathbb{R}[u_1, \ldots, u_\ell],$$

the restriction map $i_w^* : H_T^*(G/T) \longrightarrow H_T^*(\{w\})$ *is given by*

$$i_w^*(u_i) = u_i, \qquad i_w^* f(y) = w \cdot f(u),$$

for $f(y) = f(y_1, \ldots, y_\ell) \in \mathbb{R}[y_1, \ldots, y_\ell]$.

The proof of the proposition is based on two lemmas. Given a character γ of the torus T, consider the T-equivariant associated bundle $L_\gamma = L(G/T, \gamma)$ over G/T. At a fixed point w, the torus maps the fiber $(L_\gamma)_w$ linearly to itself. Hence, the fiber $(L_\gamma)_w$ is a representation of T.

Lemma 3. *At a fixed point $w \in W \subset G/T$, the fiber $(L_\gamma)_w$ is isomorphic to the representation $\mathbb{C}_{w\cdot\gamma}$ of T.*

Proof. A point in the fiber of L_γ above w is of the form $[w, v] \in G \times_T \mathbb{C}_\gamma$. For $t \in T$, since $w^{-1}tw \in T$,

$$\begin{aligned} t \cdot [w, v] &= [tw, v] = [ww^{-1}tw, v] = [w, \gamma(w^{-1}tw)v] \\ &= [w, (w \cdot \gamma)(t)v]. \end{aligned}$$

Thus, under the isomorphism

$$\begin{aligned} (L_\gamma)_w &\longrightarrow \mathbb{C} \\ [w, v] &\mapsto v, \end{aligned}$$

the torus T acts on $\mathbb{C}$ by $t \cdot v = (w \cdot \gamma)(t)v$. □

Lemma 4. *Let $j_w = (i_w)_T : \{w\}_T \longrightarrow (G/T)_T$ be the map of homotopy quotients induced by the inclusion i_w. For a character γ of T, the restriction of the line bundle $K_\gamma := (L_\gamma)_T = L(G/T, \gamma)_T$ from $(G/T)_T$ to $\{w\}_T \simeq BT$ is given by*

$$j_w^* K_\gamma \simeq S_{w\cdot\gamma}.$$

Proof. By Lemma 3 the restriction of the line bundle L_γ to the fixed point w gives rise to a commutative Cartesian diagram of T-equivariant maps

$$\begin{tikzcd} \mathbb{C}_{w\cdot\gamma} \arrow[r, hook] \arrow[d] & L_\gamma \arrow[d] \\ \{w\} \arrow[r, hook, "i_w"'] & G/T. \end{tikzcd}$$

Taking the homotopy quotients results in the Cartesian diagram

$$\begin{tikzcd} (\mathbb{C}_{w\cdot\gamma})_T \arrow[r, hook] \arrow[d] & (L_\gamma)_T \arrow[d] \\ BT \arrow[r, hook, "(i_w)_T"'] & (G/T)_T. \end{tikzcd}$$

But

$$(\mathbb{C}_{w\cdot\gamma})_T = ET \times_T \mathbb{C}_{w\cdot\gamma}$$

is precisely the line bundle $S_{w\cdot\gamma}$ over BT associated to the character $w \cdot \gamma$ of T. □

Proof of Proposition 2. Let $\pi : G/T \longrightarrow \{w\}$ be the constant map. Then $\pi \circ i_w = 1_w$, the identity map on $\{w\}$. The elements u_i in $H_T^*(G/T)$ are really $\pi^* u_i$. Hence,

$$i_w^* u_i = i_w^* \pi^* u_i = u_i.$$

In Lemma 4, take γ to be χ_i. Then

$$\begin{aligned} i_w^* y_i &= i_w^* c_1(K_{\chi_i}) = c_1(j_w^* K_{\chi_i}) = c_1(S_{w \cdot \chi_i}) \\ &= w \cdot c_1(S_{\chi_i}) \qquad \text{(by Cor. 1.2)} \\ &= w \cdot u_i. \end{aligned}$$

Since i_w^* and w are both algebra homomorphisms, for any polynomial f in ℓ variables,

$$i_w^* f(y_1, \dots, y_\ell) = w \cdot f(u_1, \cdots, u_\ell).$$

□

5. The tangent bundle of G/H

Let G be a Lie group with Lie algebra $\mathfrak{g}$, and H a closed subgroup with Lie algebra $\mathfrak{h}$.

Proposition 5 (See also [14], p. 130)**.** *The tangent bundle of G/H is diffeomorphic to the vector bundle $G \times_H (\mathfrak{g}/\mathfrak{h})$ associated to the principal H-bundle $G \longrightarrow G/H$ via the adjoint representation of H on $\mathfrak{g}/\mathfrak{h}$.*

Proof. For $g \in G$, let $\ell_g : G/H \longrightarrow G/H$ denote left multiplication by g. Then ℓ_g is a diffeomorphism and its differential

$$\ell_{g*} : T_{eH}(G/H) \longrightarrow T_{gH}(G/H)$$

is an isomorphism.

Now define

$$\varphi : G \times_H (\mathfrak{g}/\mathfrak{h}) \longrightarrow T(G/H)$$

by

$$\varphi([g, v]) = \ell_{g*}(v).$$

To show that φ is well-defined, pick another representative of $[g, v]$, say $[gh, \mathrm{Ad}(h^{-1})v]$ for some $h \in H$. Then

$$\begin{aligned} \ell_{gh*}(\mathrm{Ad}(g^{-1})v) &= \ell_{gh*}(\ell_{h^{-1}})_* r_{h*} v \\ &= \ell_{g*} \ell_{h*} (\ell_{h^{-1}})_* r_{h*} v \\ &= \ell_{g*} v, \end{aligned}$$

since right multiplication r_h on G/H is the identity map.

Since φ is clearly a surjective bundle map of two vector bundles of the same rank, it is an isomorphism. □

6. Equivariant Euler class of the normal bundle at a fixed point of G/T

In this section G is a compact connected Lie group, T a maximal torus in G, and $\mathfrak{g}$ and $\mathfrak{t}$ their respective Lie algebras. The adjoint representation of T on $\mathfrak{g}$ decomposes $\mathfrak{g}$ into a direct sum

$$\mathfrak{g} = \mathfrak{t} \oplus (\oplus_{\alpha \in \triangle^+} \mathbb{C}_\alpha),$$

where $\triangle^+$ is a choice of positive roots.

Let w be a fixed point of the left multiplication action of T on G/T, and ν_w the normal bundle of $\{w\}$ in G/T. The normal bundle at a point is simply the tangent space. Let $\ell(w)$ be the length of w and $(-1)^w := (-1)^{\ell(w)}$ the sign of w.

Proposition 6. *The equivariant Euler class of the normal bundle ν_w at a fixed point $w \in W$ is*

$$e^T(\nu_w) = w \cdot \left(\prod_{\alpha \in \triangle^+} c_1(S_\alpha) \right) = (-1)^w \prod_{\alpha \in \triangle^+} c_1(S_\alpha) \in H^*(BT).$$

Proof. By Prop. 5 the tangent bundle of G/T is the homogeneous vector bundle associated to the adjoint representation of T on $\mathfrak{g}/\mathfrak{t} = \oplus_{\alpha\in\triangle^+}\mathbb{C}_\alpha$. In the notation of Section 1,

$$T(G/T) \simeq G \times_T (\oplus_{\alpha\in\triangle^+}\mathbb{C}_\alpha) \simeq \oplus_{\alpha\in\triangle^+} L_\alpha. \tag{4}$$

By Eq. (4) and Lemma 3, at a fixed point w the normal bundle ν_w is

$$\nu_w = T_w(G/T) = \oplus_{\alpha\in\triangle^+}(L_\alpha)_w \simeq \oplus_{\alpha\in\triangle^+}\mathbb{C}_{w\cdot\alpha}.$$

It follows that the equivariant Euler class of ν_w is

$$\begin{aligned} e^T(\nu_w) &= e^T(\oplus_{\alpha\in\triangle^+}\mathbb{C}_{w\cdot\alpha}) = e(\oplus_{\alpha\in\triangle^+}(\mathbb{C}_{w\cdot\alpha})_T) \\ &= e(\oplus_{\alpha\in\triangle^+}S_{w\cdot\alpha}) && \text{(by the definition of } S_\alpha) \\ &= \prod_{\alpha\in\triangle^+} c_1(S_{w\cdot\alpha}) = \prod_{\alpha\in\triangle^+} w\cdot c_1(S_\alpha) && \text{(Cor. 1.2)} \\ &= w\cdot\left(\prod_{\alpha\in\triangle^+} c_1(S_\alpha)\right) && (w \text{ is a ring homomorphism}) \end{aligned}$$

Each simple reflection s in W_G carries exactly one positive root to a negative root. Hence,

$$s\cdot\left(\prod_{\alpha\in\triangle^+} c_1(S_\alpha)\right) = \prod_{\alpha\in\triangle^+} c_1(S_{s\cdot\alpha}) = -\prod_{\alpha\in\triangle^+} c_1(S_\alpha).$$

Since w is the product of $\ell(w)$ reflections,

$$w\cdot\left(\prod_{\alpha\in\triangle^+} c_1(S_\alpha)\right) == (-1)^{\ell(w)}\prod_{\alpha\in\triangle^+} c_1(S_\alpha).$$

□

7. EQUIVARIANT COHOMOLOGY OF G/H

As before, G is a compact connected Lie group with maximal torus T. Suppose H is a closed subgroup of G containing T. In the next few sections we indicate the modifications necessary when the homogeneous space G/T is replaced by G/H. Since there are two groups G and H containing the maximal torus T, denote the Weyl groups by W_G and W_H respectively.

The equivariant cohomology ring of G/H follows from ([8], Prop. 1(iii)), as follows. The commutative diagram

$$\begin{tikzcd} (G/H)_G \arrow[d, "\pi"'] & (G/H)_T \arrow[l, "j"'] \arrow[d, "\pi'"] \\ \mathrm{pt}_G & \mathrm{pt}_T \arrow[l] \end{tikzcd}$$

induces the commutative diagram in cohomology

$$\begin{tikzcd} H_G^*(G/H) \arrow[r, "j^*"] & H_T^*(G/H) \\ H^*(BG) \arrow[u, "\pi^*"] \arrow[r] & H^*(BT) \arrow[u, "(\pi')^*"'] \end{tikzcd},$$

and hence a ring homomorphism

$$H^*(BT)\otimes_{H^*(BG)} H_G^*(G/H) \longrightarrow H_T^*(G/H). \tag{5}$$

It is shown in ([8], Prop. 1(iii)) that Eq. (5) is an isomorphism.

For definiteness, write as in Eq. 2

$$R := H^*(BT) \simeq \mathbb{R}[u_1,\ldots,u_\ell]. \tag{6}$$

To distinguish the two copies of R in $R \otimes R^{W_H}$, we denote the second factor by $\mathbb{R}[y_1, \ldots, y_\ell]^{W_H}$. Thus,

$$H_T^*(G/H) = \frac{\mathbb{R}[u_1, \ldots, u_\ell] \otimes (\mathbb{R}[y_1, \ldots, y_\ell]^{W_H})}{J}, \tag{7}$$

where J is the ideal generated by $p(y) - p(u)$ for all W_G-invariant homogeneous polynomials $p \in R_+^{W_G}$ of positive degree in ℓ variables.

With the identifications (6) and (7), the map $j^* : H_G^*(G/H) \longrightarrow H_T^*(G/H)$ is given by

$$j^* b(u) = b(y). \tag{8}$$

8. Fixed points of T acting on G/H

The quotient space W_G/W_H can be viewed as a subset of G/H via

$$\frac{W_G}{W_H} \simeq \frac{N_G(T)}{N_H(T)} = \frac{N_G(T)}{N_G(T) \cap H} \hookrightarrow \frac{G}{H}.$$

Proposition 7. *Suppose T acts on G/H by left multiplication. Then the fixed point set F is W_G/W_H.*

Proof. The coset xH is a fixed point

$$\begin{aligned} &\text{iff} \quad txH = xH \qquad \text{for all } t \in T \\ &\text{iff} \quad x^{-1}txH = H \qquad \text{for all } t \in T \\ &\text{iff} \quad x^{-1}tx \in H \qquad \text{for all } t \in T \\ &\text{iff} \quad x^{-1}Tx \subset H. \end{aligned}$$

Since any two maximal tori in H are conjugate by an element of H, there is an element $h \in H$ such that

$$h^{-1}x^{-1}Txh = T.$$

Therefore, $xh \in N_G(T)$, and $xH = xhH$. Thus any fixed point can be represented as yH for some $y \in N_G(T)$.

Conversely, if $y \in N_G(T)$, then yH is a fixed point of the action of T on G/H. It follows that there is a surjective map

$$\begin{aligned} N_G(T) &\twoheadrightarrow F \subset G/H, \\ y &\mapsto yH \end{aligned}$$

and hence a bijection

$$\frac{N_G(T)}{H \cap N_G(T)} \simeq F.$$

□

9. The restriction to a fixed point of G/H

Theorem 8. *With $H_T^*(G/H)$ as in Eq. (7) and w a fixed point in W_G/W_H, the restriction map i_w^* in equivariant cohomology*

$$i_w^* : H_T^*(G/H) \longrightarrow H_T^*(\{w\}) = \mathbb{R}[u_1, \ldots, u_\ell]$$

is given by

$$i_w^* u_i = u_i, \qquad i_w^* f(y) = w \cdot f(u)$$

for any W_H-invariant polynomial $f(y) \in \mathbb{R}[y_1, \ldots, y_\ell]^{W_H}$.

Proof. Write $w = xH \in G/H$. The commutative diagram of T-equivariant maps

$$\begin{tikzcd} \{xT\} \arrow[r, "i_{xT}"] \arrow[d, "\sigma"'] & G/T \arrow[d, "\sigma"] \\ \{xH\} \arrow[r, "i_{xH}"'] & G/H \end{tikzcd}$$

induces a commutative diagram in equivariant cohomology

$$\begin{tikzcd}
H^*(BT) & H_T^*(G/T) \arrow[l, "i_{xT}^*"'] \\
H^*(BT) \arrow[u, "\sigma^*"] & H_T^*(G/H)\cdot \arrow[l, "i_{xH}^*"] \arrow[u, "\sigma^*"']
\end{tikzcd}$$

Since $\sigma^* : H_T^*(G/H) \longrightarrow H_T^*(G/T)$ is an injection, the restriction i_{xH}^* is given by the same formula as the restriction i_{xT}^*. The theorem then follows from Prop. 2. ☐

10. Pullback of an associated bundle

Assume that $\rho : H \longrightarrow \mathrm{Aut}(V)$ is a representation of H. By restriction, one obtains a representation of the maximal torus T.

Proposition 9. *Let $\sigma : G/T \longrightarrow G/H$ be the projection map. Under σ the associated bundle $G \times_H V$ pulls back to $G \times_T V$:*

$$\sigma^*(G \times_H V) \simeq G \times_T V.$$

Proof. Let $[g, v]_T$ and $[g, v]_H$ denote the equivalence classes of (g, v) in $G \times_T V$ and $G \times_H V$ respectively. Define $\varphi : G \times_T V \longrightarrow \sigma^*(G \times_H V)$ by

$$[g, v]_T \mapsto (gT, [g, v]_H).$$

If $(gT, [g', v']_H)$ is any element of $\sigma^*(G \times_H V)$, then $gH = g'H$. Hence, $g' = gh$ for some $h \in H$ and

$$\begin{aligned}(gT, [g', v']_H) &= (gT, [gh, v']_H)\\ &= (gT, [g, hv']_H)\\ &= \varphi([g, hv']_T),\end{aligned}$$

which shows that φ is surjective. A surjective bundle map between two vector bundles of the same rank is an isomorphism. ☐

11. Pulling back the tangent bundle of G/H to G/T

Let $\mathfrak{g}, \mathfrak{h}$, and $\mathfrak{t}$ be the Lie algebras of G, H, and T respectively. Under the adjoint representation of H the Lie algebra $\mathfrak{g}$ decomposes into a direct sum of H-modules

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}.$$

By Prop. 5, the tangent bundle of G/H is the associated bundle

$$T(G/H) \simeq G \times_H \mathfrak{m}. \tag{9}$$

Restricting the adjoint representation to the maximal torus T, the Lie algebras of H and G decompose further into a sum of T-modules

$$\mathfrak{h} = \mathfrak{t} \oplus (\oplus_{\alpha \in \triangle^+(H)} \mathbb{C}_\alpha)$$

and

$$\mathfrak{g} = \mathfrak{t} \oplus (\oplus_{\alpha \in \triangle^+(H)} \mathbb{C}_\alpha) \oplus (\oplus_{\alpha \in \triangle^+ - \triangle^+(H)} \mathbb{C}_\alpha),$$

where $\triangle^+(H)$ denotes a choice of positive roots for H, and $\triangle^+$ a choice of positive roots for G containing $\triangle^+(H)$. Hence, as a T-module,

$$\mathfrak{m} = \oplus_{\alpha \in \triangle^+ - \triangle^+(H)} \mathbb{C}_\alpha. \tag{10}$$

Proposition 10. *Under the natural projection $\sigma : G/T \longrightarrow G/H$, the tangent bundle $T(G/H)$ pulls back to a sum of associated line bundles:*

$$\sigma^* T(G/H) \simeq \oplus_{\alpha \in \triangle^+ - \triangle^+(H)} L_\alpha.$$

Proof.

$$\begin{aligned}\sigma^* T(G/H) \simeq \sigma^*(G \times_H \mathfrak{m}) \simeq G \times_T \mathfrak{m} \qquad & \text{(by (9) and Prop. 9)}\\ = G \times_T (\oplus_{\alpha \in \triangle^+ - \triangle^+(H)} \mathbb{C}_\alpha) \qquad & \text{(by (10))}\\ = \oplus_{\alpha \in \triangle^+ - \triangle^+(H)} L_\alpha \qquad & \text{(def. of } L_\alpha\text{)}\end{aligned}$$

☐

12. THE NORMAL BUNDLE AT A FIXED POINT OF G/H

At a fixed point $w = xH$ of G/H, with $x \in N_G(T)$, the action of T on G/H induces an action of T on the tangent space $T_w(G/H)$.

Proposition 11. *At a fixed point $w = xT$ of G/H, the tangent space $T_w(G/H)$ decomposes as a representation of T into*

$$T_w(G/H) \simeq \oplus_{\alpha \in \triangle^+ - \triangle^+(H)} \mathbb{C}_{w \cdot \alpha}.$$

Proof. Let $\sigma : G/T \longrightarrow G/H$ be the projection map. Then

$$\begin{aligned}
T_{xH}(G/H) &= (T(G/H))_{xH} && \text{(fiber of tangent bundle at } xH) \\
&= (\sigma^* T(G/H))_{xT} \\
&= (\oplus_{\alpha \in \triangle^+ - \triangle^+(H)} L_\alpha)_{xT} && \text{(Prop. 10)} \\
&\simeq \oplus_{\alpha \in \triangle^+ - \triangle^+(H)} \mathbb{C}_{w \cdot \alpha} && \text{(Lemma 3)}
\end{aligned}$$

□

Theorem 12. *At a fixed point $w \in W_G/W_H$ of the left action of T on G/H, the equivariant Euler class of the normal bundle ν_w is given by*

$$e^T(\nu_w) = w \cdot \left(\prod_{\alpha \in \triangle^+ - \triangle^+(H)} c_1(S_\alpha) \right).$$

Proof. The normal bundle ν_w at the point w is the tangent space $T_w(G/H)$. By Prop. 11, the equivariant Euler class of ν_w is

$$\begin{aligned}
e^T(\nu_w) &= e^T(T_w(G/H)) = e^T(\oplus_{\alpha \in \triangle^+ - \triangle^+(H)} \mathbb{C}_{w \cdot \alpha}) \\
&= \prod_{\alpha \in \triangle^+ - \triangle^+(H)} c_1^T(\mathbb{C}_{w \cdot \alpha}) = \prod c_1((\mathbb{C}_{w \cdot \alpha})_T) \\
&= \prod c_1(S_{w \cdot \alpha}) = w \cdot \left(\prod_{\alpha \in \triangle^+ - \triangle^+(H)} c_1(S_\alpha) \right).
\end{aligned}$$

□

13. EQUIVARIANT CHARACTERISTIC NUMBERS OF G/H

Suppose a torus T acts on a compact oriented manifold M. Let $\pi : M \longrightarrow \text{pt}$ be the constant map and $\pi_* : H_T^*(M) \longrightarrow H_T^*(\text{pt}) = H^*(BT)$ the push-forward or integration map in equivariant cohomology. For any class $\eta \in H_T^*(M)$, the Atiyah- Bott-Berline-Vergne (ABBV) localization formula computes $\pi_* \eta$ in terms of an integral over the fixed point set F ([3], [4]). In case the fixed points are isolated, the ABBV localization formula states that

$$\pi_* \eta = \sum_{p \in F} \frac{i_p^* \eta}{e^T(\nu_p)}.$$

For G/H with the standard torus action, let

$$f(y) \in \frac{\mathbb{R}[u_1, \ldots, u_\ell] \otimes (\mathbb{R}[y_1, \ldots, y_\ell]^{W_H})}{(p(y) - p(u) \mid p \in R_+^{W_G})}.$$

Since $y_1, \ldots, y_\ell$ are all equivariant characteristic classes, $\pi_* f(y)$ is called an *equivariant characteristic number* of G/H. With the restriction formula (Th. 8) and the Euler class formula (Th. 12),

$$\pi_* f(y) = \sum_{w \in W_G/W_H} w \cdot \left(\frac{f(u)}{\prod_{\alpha \in \triangle^+ - \triangle^+(H)} c_1(S_\alpha)} \right). \tag{11}$$

For G/T, the formula is slightly simpler. For

$$f(y) \in \frac{\mathbb{R}[u_1, \ldots, u_\ell, y_1, \ldots, y_\ell]}{(p(y) - p(u) \mid p \in R_+^{W_G})},$$

one has

$$\pi_* f(y) = \sum_{w \in W_G} w \cdot \left(\frac{f(u)}{\prod_{\alpha \in \triangle^+} c_1(S_\alpha)} \right) = \frac{\sum_{w \in W_G} (-1)^w w \cdot f(u)}{\prod_{\alpha \in \triangle^+} c_1(S_\alpha)}, \tag{12}$$

where the last equality follows from Prop. 6.

14. ORDINARY INTEGRATION AND EQUIVARIANT INTEGRATION

We state a general principle (Prop. 13), well known to the experts, relating ordinary integration and equivariant integration. It is implicit in Atiyah-Bott ([3], Section 8) and explicitly stated for the Chow ring in Edidin-Graham ([9], Prop. 5, p. 627).

Proposition 13. *Let M be a compact oriented manifold of dimension n on which a group G acts and let $\pi_* : H_G^*(M) \longrightarrow H^*(BG)$ be equivariant integration. Suppose a cohomology class $\eta \in H^n(M)$ has an equivariant extension $\tilde{\eta} \in H_G^n(M)$. Then*

$$\int_M \eta = \pi_* \tilde{\eta}. \tag{13}$$

Remark 13.1. For a torus action the right-hand side $\pi_* \tilde{\eta}$ of Eq. (13) can be computed using the ABBV localization formula in terms of the fixed point set F of T on M. In case the fixed points are isolated, this gives

$$\int_M \eta = \sum_{p \in F} \frac{i_p^* \tilde{\eta}}{e^T(\nu_p)}.$$

Proof of Prop. 13. The commutative diagram

$$\begin{tikzcd} M \arrow[r, "i"] \arrow[d, "\tau"'] & M_G \arrow[d, "\pi"] \\ \text{pt} \arrow[r, "j"'] & BG \end{tikzcd}$$

induces by the push-pull formula a commutative diagram in cohomology

$$\begin{tikzcd} H^n(M) \arrow[d, "\tau_*"'] & H_G^n(M) \arrow[l, "i^*"'] \arrow[d, "\pi_*"] \\ H^0(\text{pt}) & H^0(BG) \arrow[l, "j^*"] \end{tikzcd}. \tag{14}$$

In degree 0, the restriction $j^* : H^0(BG) \longrightarrow H^0(\text{pt})$ is an isomorphism. Hence, if η has degree n in $H^n(M)$, then by the commutative diagram (14)

$$\int_M \eta = \tau_* \eta = \tau_* i^* \tilde{\eta} = j^* \pi_* \eta = j^* \pi_* \tilde{\eta} = \pi_* \tilde{\eta}.$$

□

Since all ordinary characteristic classes of G-vector bundles have equivariant extensions, all ordinary characteristic numbers of G-vector bundles can be computed from the ABBV localization formula.

Fix a basis $\chi_1, \ldots, \chi_\ell$ of the character group $\hat{T}$ of the maximal torus T in the compact connected Lie group G. Let H be a closed subgroup containing T in G. On G/T, we have associated bundles $L_{\chi_i} := G \times_T \mathbb{C}_{\chi_i}$. Let $z_i = c_1(L_{\chi_i}) \in H^2(G/T)$. Let R be the polynomial ring $\mathbb{R}[z_1, \ldots, z_\ell]$. By Borel ([6], Th. 26.1(b)),

$$H^*(G/H) = \frac{R^{W_H}}{(R_+^{W_G})} = \frac{\mathbb{R}[z_1, \ldots, z_\ell]^{W_H}}{(R_+^{W_G})}.$$

Theorem 14. *Let* $f(z) \in \mathbb{R}[z_1, \dots, z_\ell]^{W_H}$ *be a* W_H*-invariant polynomial of degree* $\dim G/H$*, where each* z_i *has degree 2. Then the characteristic number* $\int_{G/H} f(z)$ *of* G/H *is given by*

$$\int_{G/H} f(z) = \pi_* f(y) = \sum_{w \in W_G/W_H} w \cdot \left(\frac{f(u)}{\prod_{\alpha \in \triangle^+ - \triangle^+(H)} c_1(S_\alpha)} \right).$$

Proof. Since $y_i = c_1^T(L_{\chi_i})$ is an equivariant extension of z_i, the cohomology class $f(y) \in H_T^*(G/H)$ is an equivariant extension of $f(z)$. Combining Prop. 13 and Eq. (11), the formula for the ordinary characteristic numbers of G/H follows. $\square$

15. Example: the complex Grassmannian

In this example we work out the characteristic numbers of the complex Grassmannian $G(k, n)$ of k-planes in $\mathbb{C}^n$. As a homogeneous space, $G(k, n)$ can be represented as G/H, where G is the unitary group $U(n)$ and H the closed subgroup $U(k) \times U(n-k)$.

A maximal torus containing H is

$$T = U(1) \times \cdots \times U(1) = \left\{ t = \begin{bmatrix} t_1 & & \\ & \ddots & \\ & & t_n \end{bmatrix} \;\middle|\; t_i \in U(1) \right\}.$$

A basis for the characters of T is $\chi_1, \dots, \chi_n$, with $\chi_i(t) = t_i$. A choice of positive roots for G and for H is

$$\begin{aligned} \triangle^+ &= \{\chi_i - \chi_j \mid 1 \le i < j \le n\}, \\ \triangle^+(H) &= \{\chi_i - \chi_j \mid 1 \le i < j \le k\} \cup \{\chi_i - \chi_j \mid k+1 \le i < j \le n\}. \end{aligned}$$

Therefore,

$$\triangle^+ - \triangle^+(H) = \{\chi_i - \chi_j \mid 1 \le i \le k,\ k+1 \le j \le n\}.$$

If $\alpha = \chi_i - \chi_j$, then

$$c_1(S_\alpha) = c_1(S_{\chi_i} \otimes S_{\chi_j}^*) = c_1(S_{\chi_i}) - c_1(S_{\chi_j}) = u_i - u_j. \tag{15}$$

The Weyl groups of T in G and H are

$$\begin{aligned} W_G &= S_n = \text{ the symmetric group on } n \text{ objects}, \\ W_H &= S_k \times S_{n-k}. \end{aligned}$$

Hence, the coset space W_G/W_H is in one-to-one correspondence with the set of multi-indices $I = (i_1 < \cdots < i_k)$ of length k, with $1 \le i_r \le n$ for all $r = 1, \dots, k$. Each such I has a complementary multi-index $J = (j_1 < \cdots < j_{n-k})$ with $I \cup J = \{1, \dots, n\}$.

The permutation $w \in S_n/(S_k \times S_{n-k})$ corresponding to I is

$$w(1) = i_1, \dots, w(k) = i_k, w(k+1) = j_1, \dots, w(n) = j_{n-k}. \tag{16}$$

By Eq. (7), the real equivariant cohomology ring of the Grassmannian $G(k, n)$ under the torus action is

$$H_T^*(G(k,n)) = \frac{\mathbb{R}[u_1, \dots, u_n] \otimes (\mathbb{R}[y_1, \dots, y_k, y_{k+1}, \dots, y_n]^{S_k \times S_{n-k}})}{(\prod(1+y_i) - \prod(1+u_i))} \tag{17}$$

In this formula, the notation $(p(u, y))$ means the ideal generated by the homogeneous terms of the polynomial $p(u, y)$.

Let σ_r be the r-th elementary symmetric polynomial. If S and Q are the universal sub and quotient bundles over $G(k, n)$, then

$$s_r = c_r^T(S) = \sigma_r(y_1, \dots, y_k) \quad \text{and} \quad q_r = c_r^T(Q) = \sigma_r(y_{k+1}, \dots, y_n).$$

Thus Eq. (17) can be rewritten in the form

$$H_T^*(G(k,n)) = \frac{\mathbb{R}[u_1, \dots, u_n, s_1, \dots, s_k, q_1, \dots, q_{n-k}]}{((1 + s_1 + \cdots + s_k)(1 + q_1 + \cdots + q_{n-k}) - \prod(1+u_i))}.$$

Using the relation

$$1 + \sum_{i=1}^{n-k} q_i = \frac{\prod(1+u_i)}{1 + \sum s_i}, \tag{18}$$

one can eliminate all the q_i from $H_T^*(G(k,n))$; in other words, $H_T^*(G(k,n))$ is generated as an algebra over $\mathbb{R}[u_1,\dots,u_n]$ by $s_1,\dots,s_k$ with relations given by terms of degree $> 2(n-k)$ in (18). In computing degrees, keep in mind that $\deg u_i = 2$ and $\deg s_i = \deg q_i = 2i$.

Proposition 15. *The characteristic numbers of $G(k,n)$ are*

$$\int_{G(k,n)} c_1(S)^{m_1}\dots c_k(S)^{m_k} = \sum_I \frac{\prod_{r=1}^k \sigma_r(u_{i_1},\dots,u_{i_k})^{m_r}}{\prod_{i\in I}\prod_{j\in J}(u_i-u_j)}, \tag{19}$$

where $\sum m_r = k(n-k)$ and I runs over all multi-indices $1 \le i_1 < \cdots < i_k \le n$.

Proof. In Theorem 14, take $f(y)$ to be $\prod_{r=1}^k \sigma_r(y_1,\dots,y_k)^{m_r}$. Then

$$w\cdot f(u) = \prod_{r=1}^k \sigma_r(u_{i_1},\dots,u_{i_k})^{m_r}.$$

By (15),

$$\prod_{\alpha\in\triangle^+-\triangle^+(H)} c_1(S_\alpha) = \prod_{i=1}^k\prod_{j=k+1}^n (u_i-u_j).$$

By (16),

$$w\cdot\left(\prod_{\alpha\in\triangle^+-\triangle^+(H)} c_1(S_\alpha)\right) = \prod_{i\in I}\prod_{j\in J}(u_i-u_j).$$

□

One of the surprising features of the localization formula is that although the right-hand side of Eq. (19) is apparently a sum of rational functions of $u_1,\dots,u_n$, the sum is in fact an integer.

References

[1] A. Arabia, Cohomologie T-équivariante de la variété de drapeaux d'un groupe de Kač-Moody, *Bull. Soc. math. France* 117 (1989), 129–165.

[2] E. Akyildiz and J. B. Carrell, An algebraic formula for the Gysin homomorphism from G/B to G/P, Illinois J. Math. 31 (1987), 312–320.

[3] M. Atiyah and R. Bott, The moment map and equivariant cohomology, *Topology*, 1984, 1–28.

[4] N. Berline and M. Vergne, Classes caractéristiques équivariantes. Formule de localization en cohomologie équivariante, *C. R. Acad. Sc. Paris*, Série I, t. 295 (1982), pp. 539–540.

[5] A. Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, *Ann. Math.* 57 (1953), 115–207.

[6] A. Borel et al, *Seminar on Transformation Groups*, Annals of Math. Studies 46, Princeton University Press, 1960.

[7] R. Bott, An introduction to equivariant cohomology, in *Quantum Field Theory: Perspective and Prospective* (ed. C. DeWitte-Morette), 35–56, Kluwer Academic Publishers, Dordrecht, 1999.

[8] M. Brion, Equivariant cohomology and equivariant intersection theory (Notes by Alvaro Rittatore), in NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 514, *Representation Theories and Algebraic Geometry (Montreal, PQ, 1997)*, 1–37, Kluwer Acad. Publ., Dordrecht, 1998.

[9] D. Edidin and W. Graham, Localization in equivariant intersection theory and the Bott residue formula, *Amer. J. of Math.* 120 (1998), 619–636.

[10] V. Guillemin and S. Sternberg, *Supersymmetry and Equivariant de Rham Theory*, Springer-Verlag, New York, 1999.

[11] W.-Y. Hsiang, *Cohomology of Transformation Groups*, Springer-Verlag, New York, 1975.

[12] S. K. Martin, Symplectic quotients by a nonabelian group and by its maximal torus, to appear in *Annals of Math.*

[13] A. Pedroza and L. W. Tu, Generalizaing the localization formula in equivariant cohomology, preprint.

[14] G. Segal, Equivariant K-theory, *Inst. Hautes Études Sci. Publ. Math.* 34 (1968), 129–151.

[15] L. W. Tu, The Gysin map, equivariant cohomology, and symmetrizing operators, preprint.

Department of Mathematics, Tufts University, Medford, MA 02155-7049
E-mail address: `loring.tu@tufts.edu`